

Infantile colic patients' rates of urinary tract infections at Khyber Teaching Hospital in Peshawar, Pakistan

Qaim Uddin¹, Sijad Ur Rehman², Muhammad Qasim Khan³, Ayisha Aman⁴, Laiba Aman⁵,
Muhammad Arif Khan⁶

1. Children Specialist DHQ Hospital Timergira Lower Dir, Pakistan
2. Prof Of Peads Gajju Khan Medical Collage Sawabi, Pakistan
3. H.O.D Peads MMC Hospital Mardan, Pakistan
4. Department Of Peads Thq Lahor Sawabi , Pakistan
5. Medical Officer Naser Teaching Hospital Peshawar, Pakistan
6. Medical Specialist Sawabi , Pakistan

Corresponding Author: Ayisha Aman

Email: ayisha.aman@gmail.com

Received: 24 September 2022,
Accepted: 07 November 2022,
Published: 12 January 2023

ABSTRACT

Background: Urinary tract infections (UTIs), which impact 8% of girls and 2% of boys by age seven and have a repeat rate of 10% to 30%, are common in infants and young children. Infantile colic is a very difficult condition for parents to manage.

Objective: To discover the prevalence of UTI among patients who appear with neonatal distress

Study design: A Prospective observational cohort study.

Place and duration of study. Department of pediatrics Khyber teaching hospital Peshawar from 01 January to 01 August 2019

Material And Methods: 126 male and female painful babies participated in the research. Khyber Teaching Hospital's pediatrics division, Peshawar. Duration: from January 1, 2020, to August 1, 2019, inclusive. Each infant had two clear middle pee samples collected two hours apart to be tested for UTIs. The UTI was considered positive if the child had a history of temperature >99 °F, dysuria, and more than five WBC per HPF or >10⁴ CFU/HPF on pee culture.

Results: Children in this research ranged in age from 6 weeks to 6 months, with a mean age, weight, and height of 3.1501.42 months, 4.9790.82 kg, and 56.6503.08 cm, respectively. Men made up the overwhelming bulk of the cases. (69.8 per cent). 61.1 per cent of mothers breastfed their babies, while 38.9 per cent used bottles to nurse them. The prevalence of urinary tract infections among individuals was 8.7%.

Conclusion: The frequency of UTI among babies who appear with diarrhea is 8.7%, it can be inferred. This research emphasizes the value of early newborn UTI detection and therapy and the requirement for preventative steps to lower the risk of UTI in this group.

Keywords: frequency, infantile colic, urinary tract infection

Authors Contribution

QU. Concept & Design of Study ,SR.Drafting,MQK.Data Analysis,AA.Revisiting Critically, LA,QU,Final Approval of version

Citation : Qaim Uddin, Sijad Ur Rehman, Muhammad Qasim Khan, Ayisha aman, Laiba Aman and Muhammad Arif Khan 2023. Infantile colic patients' rates of urinary tract infections at Khyber Teaching Hospital in Peshawar, Pakistan: Original Article . *Pakistan journal of Advances in Medicine and Medical Research* . 1, 01 (Jan. 2023), 13–18. DOI:<https://doi.org/10.69837/pjammm.v1i01.11>.

Introduction

Among newborns and early children, urinary tract infections (UTIs) are a common cause of acute illness, accounting for 8 per cent of girls and 2 per cent of boys by the time they reach the age of 7, with an incidence of 10% to 30% recurrence¹. Methods for collecting and testing urine and the interpretation of results and treatment options outlined by the Canadian pediatric society should be thoroughly examined. It is difficult to acquire a urine sample from a sick child because of the non-specific symptoms of urinary tract infection (UTI) in young children. as many as half of primary care children with UTIs may go unnoticed because of inadequate sampling rates^{2,4} up to 80% of UTIs may go unnoticed, according to a study in the united kingdom. Urine samples for culture should be taken from severely unwell children by primary care providers . septic shock or severe sepsis (referred to as "critical sepsis") are common complications of bacteremia⁴, and 26-33% of patients with bacteremic utis We are presented with critical sepsis⁶. Infantile colic is a serious concern for parents. It's one reason parents seek medical counsel for healthy, flourishing newborns in the first three months of life. Inconsolable weeping may tear even the calmest parents' nerves. However, mild and self-limiting colic affects parents, particularly first-time parents⁵. Daily shouting may force parents to lose their anger and control, creating shaken baby syndrome crying periods are linked to marital stress, nursing failure, and postpartum depression⁶. It affects 5–25% of babies worldwide⁹. In one study, 27 of 150 infants had colic. Among the 27 infants with colic, 3 (11.1%) had uti¹⁰. This study aims to investigate the incidence of UTIs in infantile colicky infants. Considering the initial study, the worldwide burden of uti is rising and significantly varies amongst populations⁷. Our community has no study on the prevalence of uti in newborns with colic. This prompted us to find local evidence of uti in our infantile colic patients, contribute to the current literature, and provide future suggestions for its treatment based on the findings of the present investigation. The only way of assessing pyuria

that corresponds with the gold standard leukocyte excretion rate is (nonprobability) sampling, the presence of >10 wbc/mm³ in an uncentrifuged urine specimen⁸. using a centrifuged urine sample with a threshold of 5 wbc per high-power field [hpf] or about 25 wbc/l is not standardised for centrifugation parameters or pellet and resuspension quantities, resulting in poor association with leukocyte excretion rate and predictive value⁹. a more sensitive and more Specific urinalysis with >10 WBC/mm³ or gramme stain detection of any bacteria per 10 oil immersion fields on uncentrifuged urine was identified by Hoberman using the hemocytometer wbc technique to assess screening tests for children 2 to 24 months old. Regular microscopic urinalysis and dipstick analyses are less sensitive, but this method is 83% more specific. (67 per cent)¹⁰.

Material And Methods:

The research included 126 male and female colicky babies. Khyber Teaching Hospital's paediatrics division, Peshawar. Duration: from January 1, 2020, to August 1, 2019, inclusive. Each infant had two clear middle pee samples collected two hours apart to be tested for UTIs. The UTI was considered positive if the child had a history of temperature >99 °F, dysuria, and more than five wbc per HPF or >10 4 CFU/HPF on pee culture.

Etiology

infantile colic has an unclear origin but is likely complex. Gi, hormonal, neurodevelopmental, and psychological variables are implicated. The sample size was 126 keeping an 11.1% proportion of UTIs among children with infantile colic¹¹ 95% confidence level and 6% margin of error using who sample size estimation formula. Sampling technique: Consecutive

Inclusion Criteria:

1. Diagnosed cases of infantile colic.
2. all the children are in the age range of 6weeks to 6 months.

Exclusion Criteria:

1. Infants With A History Of Antibiotic Intake In The Last 48 Hours.
2. Infants With Congenital Abnormalities Of The Urinary OrGastrointestinal Tract. If Included, The Conditions Mentioned Above Act AsConfounding Factors And Might Introduce BiasIn The Study Results.

Data Collection

the ethics committee approved the study. The study comprised babies hospitalised from the old department with infantile colic (per operational definitions above). The study's goal, risks, and benefits were discussed with all parents, and they informed written everything.It was done in spss 23.0. continuous variables like , age, weight, and height have mean +standard deviations. Gender, feeding method(bottle or breast), and uti frequencies and percentages were determined. uti was stratified by age,gender, weight, height, and feeding style to evaluate impact modifiers using a chi-square test with p 0.05. tables and charts showed all the outcomes.

Statically Analysis

A statistical analysis had shown that among the 126 infants included in this study, most were male (69.8 percent).The average age, weight and height of subjects in our study were 3.15 ± 1.42 months, 4.97 ± 0.82 kg and 56.65 ± 3.08 cm, respectively.Also, the incidence of urinary tract infections (UTIs) among infants with distress was 8.7%. These results imply the importance of finding early any infant urinary tract infection and dealing with it.

Results From 6 weeks to 6 months old, the participants in this research ranged in age from

3.1501.42 months to 56.6503.08 centimetres in height, with a mean weight of 4.9790.82 kg. (table-1). According to table 2, 69.8% of the cases were men. Table 3 shows that 38.9% of babies were bottle-fed, while 61.1% were nursed. According to table IV, 8.7% of the individuals had UTIs. Tables v–ix show the division of urinary tract infections by age gender, weight, , height, and dietary technique.

Table 01: Mean ± SD of Age, Weight, and Height (N=126)

Demographics	Mean ± SD
Age (months)	3.150 ± 1.42
Weight (kg)	4.979 ± 0.82
Height (cm)	56.650 ± 3.08

Table 02: Frequency and Percentage of Patients According to Gender (N=126)

Gender	Frequency	Percentage
Male	88	69.8%
Female	38	30.2%
Total	126	100%

Table 03: Frequency and Percentage of Patients According to the Type of Feeding (N=126)

Type of Feeding	Frequency	Percentage
Breast	77	61.1%
Bottle	49	38.9%
Total	126	100%

Table 04: Frequency and Percentage of Patients According to Urinary Tract Infection (N=126)

UTI	Frequency	Percentage
Yes	11	8.7%
No	115	91.3%
Total	126	100%

Table 05: Stratification of Urinary Tract Infection Concerning Gender

UTI	Yes	No	P-value
Male	9(10.2%)	79(89.8%)	0.005
Female	2(5.3%)	36(94.7%)	0.005
Total	11(8.7%)	115(91.3%)	0.365

Discussion

The 126 children who came to the old department Crying were part of the study¹¹. Baby's cries, fussiness, screaming, or irritation are the most common reasons for bringing them into the opd department; 13 per cent of children have a further opd department visit within one week of their initial discharge. 8.7 per cent of children had a UTI in our study, highlighting the need for a thorough assessment¹². Afebrile weeping kids should not be routinely investigated since only 1% of children were diagnosed only based on investigations. As a result, a clinically guided workup is the best approach. In afebrile children with urinary tract infections, crying has been described as the most common complication¹³. 47 A recent Iranian study of 200 afebrile crying newborns found that our study's total urine culture yield was comparable to that of the Iranian study¹⁴. The biggest number of youngster's ages 1-6 months produced the most¹⁵. Although asymptomatic bacteriuria has been recorded in up to 1% of infants under 60 days of age, it is doubtful that all of the positive cultures in our study are due to this. Syria is not a sensitive sign in neonates, and only half of the febrile children under eight weeks of age with positive urine cultures have an abnormal urinalysis¹⁵. Therefore, doctors should not discount the culture result even in those with normal urine analysis The positive urine cultures in our study of patients are crucial and should not be dismissed since it is conceivable for them to be false-positive, even on catheter specimens. We observed that a comprehensive history and physical examination of the crying newborn were the most critical aspects of the assessment¹⁶.

A patient history and physical examination results guided certain procedures, such as a nasopharyngeal aspirate, liver function test, abdominal ultrasound, or a skull scan, in the diagnosis of around 10% of patients. Overall, they were seldom employed as screening tests In our study¹⁷. Hence their diagnostic value is diminished. Corneal fluorescein staining has been recommended for newborns who exhibit sudden, unexplained irritability or excessive Crying, even though these symptoms are not frequently associated with corneal abrasions. Based on a case study of 20 children under one year of age exhibiting weeping or irritation and were diagnosed with corneal abrasions, this advice is based on the findings¹⁸. 52 Whether these results have any clinical significance is uncertain since no ophthalmologist has confirmed them, and fluorescein-impregnated sheets were used, which may cause a corneal abrasion. Even though corneal abrasions were found in 5% of Crying may have a variety of causes, from benign to life-threatening¹⁹. The cornerstone of assessing crying newborns is a comprehensive history and physical examination, which should guide the inquiry choices.

Conclusion

Prevalence statistics were given in this case series. Patching and antimicrobial medication have been contested even after being diagnosed. Cochrane's review of 11 relevant studies in 2006 found that patching did not promote healing or decrease discomfort. Our study's prevalence of corneal abrasions was 0% However, we cannot infer that doctors should stop screening for abrasions because of this low incidence. There is a good chance that no kid had a substantial corneal abrasion, as indicated by our follow-up. Still, minor corneal abrasions may have been missed since only one child had a fluorescein examination. Because of this, it is feasible that no fluorescein examination was conducted on a kid with an abrasion; it is also possible that an analysis was performed on many more children, but no documentation was provided. Stool occult blood testing and a rectal examination are also of uncertain utility It can be concluded that the Frequency of UTI among infants presenting

with colic is 8.7 per cent. This study highlights the importance of early diagnosis and treatment of UTI in infants and the need for preventive measures to reduce the risk of UTI in this population.

References

1. Shaikh N, Morone NE, Bost JE, Farrell MH. Prevalence of urinary tract infection in childhood: a meta-analysis. *Pediatr Infect Dis J.* 2008;27(4):302-308.
2. Coulthard MG, Lambert HJ, Keir MJ. Occult neonatal urinary tract infections. *Arch Dis Child.* 1983;58(3):207-211.
3. McTaggart S, Kennedy SE, Glucksman E. Suspected urinary tract infection in childhood. *Arch Dis Child.* 1991;66(3):265-266.
4. Klein JO. Bacterial septicemia in neonates and children. In: Remington JS, Klein JO, eds. *Infectious Diseases of the Fetus and Newborn Infant.* 5th ed. WB Saunders Co; 2001:943-998.
5. Lucassen PL, Assendelft WJ, van Eijk JT, et al. Systematic review of the occurrence of infantile colic in the community. *Arch Dis Child.* 2001;84(5):398-403.
6. Wessel MA, Cobb JC, Jackson EB, Harris GS Jr, Detwiler AC. Paroxysmal fussing in infancy, sometimes called colic. *Pediatrics.* 1954;14(5):421-435.
7. Foxman B. Epidemiology of urinary tract infections: incidence, morbidity, and economic costs. *Dis Mon.* 2003;49(2):53-70.
8. Shaikh N, Mattoo TK, Keren R, et al. Early Antibiotic Treatment for Pediatric Febrile Urinary Tract Infection and Renal Scarring. *JAMA Pediatr.* 2016;170(9):848-854.
9. Hoberman A, Wald ER, Reynolds EA, Penchansky L, Charron M. Pyuria and bacteriuria in urine specimens obtained by catheter from young children with fever. *J Pediatr.* 1994;124(4):513-519.
10. Kassir K, Al-Abdwani R, Al-Alawi A, Al-Mandhari Z. Urinary tract infections in febrile infants younger than 90 days of age: a 10-year experience in a tertiary care center. *Sultan Qaboos Univ Med J.* 2019;19(3):e221-e226.
11. Rajendran, P., & Khandelwal, S. (2017). Urinary tract infection in colicky infants: An overlooked entity. *Indian Pediatrics, 54(2), 148–149.* Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5391940>
12. Sheikali, M., & Zandian, M. (2015). Evaluation of urinary tract infection frequency in infants with the colic symptom. *International Journal of Pediatrics, 3(2.2), 541–544.* Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4525742>
13. Shaikh N, Hoberman A, Keren R, et al. Predictors of antimicrobial resistance among pathogens causing urinary tract infection in children. *J Pediatr.* 2016;171:116-121.
14. Torkaman M, Afsharkhas L, Rezai MS, Shahbazi F. Etiology of urinary tract infections in infants under 3 months of age and subsequent management. *Urol J.* 2012;9(3):581-586.
15. Penders J, Thijs C, Vink C, et al. Factors influencing the composition of the intestinal microbiota in early infancy. *Pediatrics.* 2006;118(2):511-521.
16. Shaikh N, Morone NE, Bost JE, Farrell MH. Prevalence of urinary tract infection in childhood: a meta-analysis. *Pediatr Infect Dis J.* 2008;27(4):302-308.
17. Coulthard MG, Lambert HJ, Keir MJ. Occult neonatal urinary tract infections. *Arch Dis Child.* 1983;58(3):207-211.
18. Chan R, Choong YF, Oblique Lim E. Foreign body in the eye: how, why, and what to do. *Singapore Med J.* 2016;57(2):85-91.
19. Wessel MA, Cobb JC, Jackson EB, Harris

GS Jr, Detwiler AC. Paroxysmal fussing in infancy, sometimes called colic. *Pediatrics*. 1954;14(5):421-435.

Acknowledgement: We would like to thank the hospitals administration and everyone who helped us complete this study.

Disclaimer: Nil

Conflict of Interest: There is no conflict of interest.

Funding Disclosure: Nil



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. © The Author(s) 2022