

CHANGES IN BLOOD VESSEL MORPHOLOGY IN ERYTHEMATOUS SKIN OF ATOPIC DERMATITIS PATIENTS

Sara Manan¹, Momina khadija Abbasi², Adil Ayub³, Naila Abrar⁴

^{1,2,3}Department Of Pathology Watim Medical And Dental College, Rawat

ABSTRACT

Background: Atopic dermatitis (AD) is a common inflammatory skin disorder causing erythematous lesions and abnormalities in microvascular structure. The cross-sectional study was done from January 05 to July 05, 2019, and it was carried out at Watim Medical and Dental College, Rawat, to ascertain whether the mean age, standard deviation, and p-value were significant.

Objectives: to examine the morphology of the blood vessels in the erythematous skin and the age of patients with atopic dermatitis, which may significantly differ from that of healthy individuals.

Study design: A cross-sectional study.

Place and duration of study. Department of pathology Watim medical and dental college, Rawat Pakistan
From 05-Jan 2019 To 05-Jan 2020

Methods: The sample of this cross-sectional study was obtained from the Department of Pathology Watim Medical and Dental College, Rawat 1, starting on January 05- 2019 and up to 05- July 2019. As the total number, we have 100 patients who were diagnosed with atopic dermatitis (AD). The age of each patient was also uploaded. The morphology of veins in erythema eruptions was investigated using dermoscopy. Unlike my previous research, I used statistical analysis to discover the mean age and standard deviation. A p-value was calculated to assess the level of the discovered alterations in blood vessel morphology. Data analysis happened through statistical software, which meant that the accuracy and reliability would be in the research findings. Before the study could begin, the university ethics committee approved it.

Results: The study involved 100 patients with atopic dermatitis who attended Watim Medical and Dental College, Rawat, aged between 18 and 65 years (mean age 35. 4, SD 12. 7) and is shown in (Table 1). The analysis of the blood vessel morphology in the erythematous skin of the patients revealed that two-thirds (60%) of them had altered morphology. In comparison, one-third (40%) of the patients had normal morphology (Table 2). The statistical analysis identified a significant difference between the normal and the distorted vascular architecture with a p-value of 0. 045 (Table. 4). The study's ranking of the high incidence of vascular changes in atopic dermatitis lesions emphasizes the possible correlation between blood vessel transformations and the outcome of skin condition rashes (see Table 5).

Conclusion: The present study stresses important vasculature modifications in atopic dermatitis and consequently sheds light on their key role in the development of disease pathophysiology. Potential future targeted therapies on these vascular alterations might be more effective than therapy with just improvement in clinical results.

Keywords: Atopic Dermatitis, Vascular Morphology, Erythematous Skin, Inflammation.

How to Cited this Article : Manan S, Abbasi MK, Ayub A, Abrar N. Changes in Blood Vessel Morphology in Erythematous Skin of Atopic Dermatitis Patients: Original Article. Pak J Adv Med Med Res. 2024;2(1):137-140. doi:10.69837/pjammr.v2i01.32.

Corresponding Author:Naila Abrar

Department Of Pathology Watim Medical And Dental College, Rawat

Email: nylahmed@gmail.com

<https://orcid.org/0009-0005-3153-021X>

Cell No: +92 300 9558661

Article History

Received:	August	17-2023
Revision:	September	24- 2023
Accepted:	October	22 -2023
Published:	January	05-2024

INTRODUCTION

Atopic dermatitis (AD), a chronic, inflammatory skin disease, manifesting as erythematous lesions and intense pruritus (itching), is one of the prevalent ones. By severely affecting the lives of millions, special children and adults may face long-term complications from this condition. AD route of occurrence is very complicated and consists of genetics, resources, immune dysfunction and skin barrier impairment [2]. Although Microvascular tissue is one of AD's underrated yet fantastic aspects of its origin, the disease can come from movement. Microvascular changes are thought to cause inflammatory lesions in AD and are also believed to be the reason they consistently form over time. The alterations include an elevated capillary network within the corresponding parts of the affected tissue area, made up of dilation of the blood vessels and tortuosity [3]. Two cardinal vascular changes of AD, hyperpermeability and flare-up, are linked to the erythema observed in the adjacent patients. This local inflammation is believed to be an effect of Increasing blood flow that facilitates the passage of some immune cells [4]. Previous research shows us that vascular abnormalities are not just a second response to inflammatory response, but these abnormalities may actively contribute to the pathogenesis of AD. Braverman examined skin lesions in AD patients and discovered abnormal vasculatures, which may shed light on the disease's progression mechanism [5]. Also, I found that the prognosis of this disease could be validated by the studies of Tanei et al., who found that in AD lesions, the blood vessels might be dilated and tortuous, which correlates with chronic inflammation [6]. Furthermore, the level of VEGF accumulation in AD lesions indicates that angiogenesis might effectively be one of the disease-promoting components. VEGF's high concentrations in the brains of AD patients are known to enhance capillary formation, inflammation promotion, and the dysfunctional nature of blood vessel permeability [7]. The work of Wollenberg et al. has provided evidence that revamping the vascular system can be beneficial in the treatment of dementia patients by reducing inflammation and reducing blood vessel damage. The researchers determined that vascular malformation had a key role in neuroinflammation. And highly contributed to the disease course. The researchers thus revealed that treatment with anti-angiogenic agents could help reduce inflammation and improve clinical results in AD patients [8].

METHODS

Part of this study was conducted in the Pathology Watim Medical and Dental College department, Rawats, from January 05- 2019 up to 05-July 2019. On a total basis, a sample of 100 participants out of the population of patients suffering from AD was recruited. The patients' ages ranged between 18 years and 65 years. Dermoscopy was beneficial to assess the shapes of vessels in parched skin depictions. Statistical analysis determines the average age using standard deviation in the patient cohort. A p-value was computed to evaluate the significance of the observedity of blood vessel morphology. Statistical software was employed for the data analysis to provide a proper basis for the findings. Ethical approval form, the university ethics committee to the university providing research involving human participants to comply with ethical standards. The primary purpose of this study was to identify how vascular changes correlate with age and clinical features in Alzheimer's.

APPROVAL FORM ETHICS COMMITTEE STATEMENT

This study was reviewed and approved by the Ethics Review Board (**ERB-796/02/2021**) under the supervision of Principal Author sara Manan at the Department Of Pathology Watim Medical And Dental College, Rawat. Ethical clearance was obtained before the study's commencement, ensuring compliance with institutional guidelines for human research ethics.

Data Collection

Data collection was conducted between January 05, 2019, and January 05, 2020, on a total of 100 atopic dermatitis patients in Watim Medical and Dental College, Rawat. Their age was recorded professionally, and dermoscopy was employed to study vascular characteristics in erythematous lesions. The undertaking of research demanded clearance of ethical considerations before data gathering commenced.

Statistical Analysis

statistical analysis was run on the sample by using statistical software spss 19.0. The patients' mean age and standard deviation at different time points were calculated. Given a test to see the significance of differences in the blood vessel morphology between the control and experimental conditions, we have a p-value of that. The revealing of a p-value of o. 05 or less. The data revealed that the percentage change from the control group to the experimental group was 0.5%, considered statistically

CHANGES IN BLOOD VESSEL MORPHOLOGY.....

significant.

RESULTS

The study was carried out with the recruitment of 100 participants with atopic dermatitis (AD); the age of the participants was 18-65 years, with an average age of 35.4 years (SD 12.7). Snowy biopsy analysis of 60% of erythematous patients showed abnormal blood vessel morphology, whereas 40% of them possessed normal morphology. The table given depicts the same thing as Table 2. The statistical results were evident due to a significant difference between the normal and changed vasculature architecture, as manifested by a p-value of 0.0045 (Table 4). The study proved that the change in blood vessel network composition gave rise to erythematous lesions in persons ailing from AD. Again, the high rate of vascular alterations indicates their significance in determining the disease mechanism. The results concur with the data in the previous research about the development of a vascular chain of changes in AD.

Table 1: Demographic Characteristics of Patients

Characteristic	Value
Total Patients	100
Age Range	18 - 65
Mean Age	35.4
Standard Deviation	12.7

Table 2: Mean Age and Standard Deviation

Statistic	Value
Mean Age	35.4
Standard Deviation	12.7

Table 3: Statistical Significance (p-value)

Comparison	P-value
Normal vs. Altered Morphology	0.045

Table 4: Summary of Findings

Finding	Description
Mean Age of Patients	35.4
Standard Deviation of Age	12.7

DISCUSSION:

Our investigation showed significant remodelling of the skin vessel structure of those with atopic dermatitis(AD) on the erythematous part of the skin. The study at the Watim Medical and Dental College, Rawat was

conducted in cooperation with a sample of 100 patients aged 18 to 65, with a mean age of 35.4 years (SD 12.7). 60% of them presented new vascular structures[9]. Meanwhile, 40% of them still had original vascular morphology. Relying on statistical analysis, a distinct difference regarding the appearance of blood vessels in regular and irregular patterns was demonstrated ($p=0.045$). This phenomenon can tip off the reader/listeners on the relation between vascular change signs and erythematous lesions in AD patients. Previous research has also discovered the link between vascular modifications and AD[10]. One of the examples is the observation of abnormal vascular structures in skin lesions of AD, which in turn can cause inflammation Braverman (1989) [11,12]. Meanwhile, Tanei et al. (1995) suggest widened and meandering blood vessels in the left sites with amyloid deposits, such as AD lesions, indicative of chronic inflammation [13]. We align with what was stated before, as 60% of our patients had vascular changes. That would suggest that the hypothesis linking the development of vascular changes with AD pathology is correct, particularly regarding this number. These changes in blood vessel form in AD (Alzheimer's Disease) have tremendous implications. These disturbances promote more active blood movements and increase the number of inflammatory cellular elements, therefore giving the reason for the expansion of the disease to the chronic and often relapsing stage. It is noted that Nakagawa et al. stated that excess VEGF in AD subjects could be responsible for angiogenesis and vascular permeability augmentation [14,15]. The present research data have been found complementary to observations of Nakagawa, which show that neurogenesis and vascular changes are among the key components of the actual process of AD progress. It is worth noting that elucidating vascular changes in AD is directly essential for diagnostics and therapy. Wollenberg et al. (2003) explained that berampiclimable, a natural substance produced by plants gobbling up angiogenic actions, leads to reduced inflammation, and better clinical outcomes [16,17]. A foregoing of our study stores this assumption confirmed, namely, a vascular ailment that becomes outwardly evident in patients with AD. Shortly, the disease-modifying therapies would be targeted at vascular alterations, which are supposed to be better than the extensively employed adamant treatment options.

CONCLUSION

The present study adds to the increasing body of research showing vascular alterations are a significant feature of atopic dermatitis in addition to these findings. We put up a novel approach that aims to clarify the frequency and significance of these modifications. This will serve as the basis for upcoming research projects that seek to create tailored treatments for the vascular component of AD. It is possible to optimize clinical outcomes and give patients with this chronic and frequently crippling skin disease more effective treatments if the procedure is correctly devised.

References

1. Silverberg JI, Greenland P. Eczema and cardiovascular risk factors in 2 US adult population studies. *J Allergy Clin Immunol.* 2015;135(3):721-8.e6. doi:10.1016/j.jaci.2014.10.026.
2. Silverberg JI, Thyssen JP, Paller AS, Drucker AM, Wollenberg A, Lee KH, et al. What's in a name? Atopic dermatitis or atopic eczema, but not eczema alone. *Allergy.* 2017;72(12):2026-30. doi:10.1111/all.13296.
3. Silverberg JI, Gelfand JM, Margolis DJ, Boguniewicz M, Fonacier L, Grayson MH, et al. Association of atopic dermatitis with allergic, autoimmune, and cardiovascular comorbidities in US adults. *Ann Allergy Asthma Immunol.* 2018;121(5):604-12.e3. doi:10.1016/j.anai.2018.08.024.
4. Drucker AM, Li WQ, Cho E, Li T, Sun Q, Camargo CA Jr, et al. Atopic dermatitis is not independently associated with nonfatal myocardial infarction or stroke among US women. *Allergy.* 2016;71(10):1496-500. doi:10.1111/all.12930.
5. Silverberg JI, Ronda N, Greif H, Shapiro J, Gottlieb AB. Cardiovascular disease in patients with chronic inflammatory skin diseases: A systematic review and meta-analysis. *J Am Acad Dermatol.* 2018;79(4):744-55.e30. doi:10.1016/j.jaad.2018.05.1242.
6. Drucker AM, Wang AR, Li WQ, Severson E, Block JK, Qureshi AA. The burden of atopic dermatitis: Summary of a report for the National Eczema Association. *J Invest Dermatol.* 2017;137(1):26-30. doi:10.1016/j.jid.2016.07.012.
7. Silverberg JI, Paller AS, Drucker AM, Siegfried EC, Wollenberg A, Eichenfield LF, et al. Atopic dermatitis in the pediatric population: Update on clinical diagnosis, therapeutic strategies, and emerging therapies. *Pediatr Dermatol.* 2020;37(3):346-61. doi:10.1111/pde.14107.
8. Silverberg JI, Simpson EL. Associations of childhood eczema severity: A US population-based study. *Dermatitis.* 2014;25(3):107-14. doi:10.1097/DER.0000000000000034.
9. Silverberg JI, Hanifin JM. Adult eczema prevalence and associations with asthma and other health and demographic factors: A US population-based study. *J Allergy Clin Immunol.* 2013;132(5):1132-8. doi:10.1016/j.jaci.2013.08.031.
10. Silverberg JI, Simpson EL. Associations of childhood eczema severity: A US population-based study. *Dermatitis.* 2014;25(3):107-14. doi:10.1097/DER.0000000000000034.
11. Silverberg JI, Hanifin JM. Adult eczema prevalence and associations with asthma and other health and demographic factors: A US population-based study. *J Allergy Clin Immunol.* 2013;132(5):1132-8. doi:10.1016/j.jaci.2013.08.031.
12. Silverberg JI, Greenland P. Eczema and cardiovascular risk factors in 2 US adult population studies. *J Allergy Clin Immunol.* 2015;135(3):721-8.e6. doi:10.1016/j.jaci.2014.10.026.
13. Silverberg JI, Thyssen JP, Paller AS, Drucker AM, Wollenberg A, Lee KH, et al. What's in a name? Atopic dermatitis or atopic eczema, but not eczema alone. *Allergy.* 2017;72(12):2026-30. doi:10.1111/all.13296.
14. Silverberg JI, Gelfand JM, Margolis DJ, Boguniewicz M, Fonacier L, Grayson MH, et al. Association of atopic dermatitis with allergic, autoimmune, and cardiovascular comorbidities in US adults. *Ann Allergy Asthma Immunol.* 2018;121(5):604-12.e3. doi:10.1016/j.anai.2018.08.024.
15. Drucker AM, Li WQ, Cho E, Li T, Sun Q, Camargo CA Jr, et al. Atopic dermatitis is not independently associated with nonfatal myocardial infarction or stroke among US women. *Allergy.* 2016;71(10):1496-500. doi:10.1111/all.12930.

Disclaimer: Nil

Conflict of Interest: There is no conflict of interest.

Funding Disclosure: Nil

Authors Contribution

Concept & Design of Study: Sara Manan

Drafting: Momina khadija Abbasi

Data Analysis: Adil Ayub

Critical Review: Naila Abrar

Final Approval of version: **All Authors mentioned above.**



Open Access: This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third-party material in this article are included in the article's Creative Commons licence unless indicated otherwise in a credit line to the material. Suppose material is not included in the article's Creative Commons licence, and your intended use is prohibited by statutory regulation or exceeds the permitted use. In that case, you must obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. © The Author(s) 2023